



	Turf: Seed	Turf: New Sod	Turf: Established	Plantings: Ball & Burlap or Container	Plantings: Soil Injection	Transplants	Compost Tea
<b>MycoApply® Ultrafine Endo (Suspendable Powder)</b>	Mix with seed at a rate of 0.37 to 0.52 ounce per 1,000 square feet of seed (11.3 - 15.9 grams per 100 square meters of seed).	Follow label for dilution directions. Apply as a liquid suspension drench to bare soil before sod is installed at a rate of 0.37 to 0.52 ounce per 1,000 square feet (11.3 - 15.9 grams per 100 square meters). Applied at a volume of 1 - 2 gallons per 1,000 ft <sup>2</sup> (4.07 - 8.15 liters per 100 m <sup>2</sup> ).	Follow label for dilution directions. Apply as a liquid suspension drench to aerated or dethatched turf at a rate of 0.52 ounce per 1,000 square feet (15.9 grams per 100 square meters). Applied at a volume of 2 gallons per 1,000 ft <sup>2</sup> (8.15 liters per 100 m <sup>2</sup> ). Irrigate well to ensure movement into the root zone.	Incorporate 0.5 ounce with 2 cubic feet of backfill soil (2.5 grams per 10 liters) or make a water suspension by mixing 2 ounces in 5 gallons of water (3 grams per liter) and apply sufficient suspension to moisten the sides of the root ball and the backfill soil.	Prepare a suspension using 1 pound per 40 gallons of water (3 grams per liter) and inject into multiple holes near the dripline of the trees or shrubs.	Dip plugs in a suspension of 2 ounces per 5 gallons of water (3 grams per liter) for 10-15 seconds to obtain good contact with substrate. For bare-rooted transplants, a spreader sticker may be added to the suspension, and the roots should be thoroughly sprayed with the suspension.	Add 1 pound per 40 gallons of tea (3 grams per liter) prior to application as a soil drench. Apply tea at a normal rate. Irrigate well after application to ensure propagule movement into the root zone.

	Turf: Seed	Turf: New Sod	Turf: Established	Plantings: Ball & Burlap or Container	Plantings: Broadcast	Transplants
<b>MycoApply® Endo (Granular)</b>	Mix with seed at a rate of 1 ounce per 1,000 square feet of seed (30.5 grams per 100 square meters of seed).	Broadcast over bare soil before sod is installed at a rate of 0.8 - 1.12 ounces per 1,000 square feet (24.5 - 34.2 grams per 100 square meters).	Apply as a broadcast during aerification of turf at a rate of 0.8 - 1.12 ounces per 1,000 square feet (24.5 - 34.2 grams per 100 square meters). Irrigate well to ensure movement into the root zone.	Incorporate 1 ounce per 2 cubic feet (5 grams per 10 liters) of backfill soil / growing media / potting mix.	Apply 1.4 ounces per 1,000 square feet (42.7 grams per 100 square meters).	Incorporate 1 ounce per cubic foot (1 gram per liter) of backfill soil / growing media / potting soil.

*The product can be blended with fertilizer, if desired.*

*If multiple applications are made in a year, then the rate can be reduced by 40% in subsequent applications.*

*MycoApply Products are compatible with a variety of fungicides. Check our website or contact us for a full compatibility list before applying a fungicide.*



	Plantings: Ball & Burlap or Container	Plantings: Soil Injection	Transplants	Compost Tea
<b>MycoApply® Ultrafine Endo/Ecto</b> (Suspendable Powder)	Incorporate 0.5 ounce with 2 cubic feet of backfill soil (3 grams per 10 liters) or make a water suspension by mixing 2 ounces in 5 gallons of water (3 grams per liter) and apply sufficient suspension to moisten the sides of the root ball and the backfill soil.	Prepare a suspension using 1 pound per 40 gallons of water (3 grams per liter) and inject into multiple holes near the dripline of the trees or shrubs.	Dip plugs in a suspension of 2 ounces per 5 gallons of water (3 g per liter) for 10 - 15 seconds to obtain good contact with substrate. For bare-rooted transplants, a spreader sticker may be added to the suspension, and the roots should be thoroughly sprayed with the suspension.	Add 1 pound per 40 gallons of tea (3 grams per liter) prior to application as a soil drench. Apply tea at a normal rate. Irrigate well after application to ensure movement into the root zone.
<b>MycoApply® Soluble MAXX</b> (Suspendable Powder)	Incorporate 0.5 ounce with 2 cubic feet of backfill soil (3 grams per 10 liters) or make a water suspension by mixing 2 ounces in 5 gallons of water (3 grams per liter) and apply sufficient suspension to saturate the sides of the root ball and the backfill soil.	Prepare a suspension using 1 pound per 40 gallons of water (3 grams per liter) and inject into multiple holes near the dripline of the trees or shrubs.	Dip plugs in a suspension of 2 ounces per 5 gallons of water (3 g per liter) for 10 - 15 seconds to obtain good contact with substrate. For bare-rooted transplants, a spreader sticker may be added to the suspension, and the roots should be thoroughly sprayed with the suspension.	Add 1 pound per 40 gallons of tea (3 grams per liter) prior to application as a soil drench. Apply tea at a normal rate. Irrigate well after application to ensure movement into the root zone.
	Plantings: Ball & Burlap or Container	Plantings: Broadcast	Transplants	
<b>MycoApply® Endo/Ecto</b> (Granular)	Incorporate 1 ounce per 2 cubic feet (5 grams per 10 liters) of backfill soil / growing media / potting mix.	Apply 1.4 ounces per 1,000 square feet (42.7 grams per 100 square meters).	Incorporate 1 ounce per cubic foot (1 gram per liter) of backfill soil / growing media / potting soil.	
	Turf: Seed	Turf: New Sod	Turf: Established	
<b>MycoApply® Injector Endo &amp; Injector Endo Organic</b> (Concentrated Suspendable Powder)	Follow label for dilution directions. Apply as a liquid suspension drench to bare soil during seeding at a rate of 0.85 - 1.2* gallons of drench per 1,000 square feet (3.5 - 4.9* liters per 100 square meters). Application must be made before seed covering is applied. Can be applied via a preblended mix tank or a 1:100 injector using a sprayer or dribbler. <i>*Application rate depends on application equipment, application site (foliar vs. direct to root zone) and turf type (density of turf)</i>	Follow label for dilution directions. Apply as a liquid suspension drench to bare soil before sod is installed at a rate of 0.85 - 1.2* gallons of drench per 1,000 square feet (3.5 - 4.9* liters per 100 square meters). Can be applied via a preblended mix tank or a 1:100 injector using a sprayer or dribbler. <i>*Application rate depends on application equipment, application site (foliar vs. direct to root zone) and turf type (density of turf)</i>	Follow label for dilution directions. Apply as a liquid suspension drench to aerated or dethatched turf at a rate of 0.85 - 1.2* gallons of drench per 1,000 square feet (3.5 - 4.9* liters per 100 square meters). Can be applied via a preblended mix tank or a 1:100 injector using a sprayer or a dribbler behind a tractor. Irrigate well to ensure movement into the root zone. <i>*Application rate depends on application equipment, application site (foliar vs. direct to root zone) and turf type (density of turf)</i>	

*The product can be blended with fertilizer, if desired.*

*If multiple applications are made in a year, then the rate can be reduced by 40% in subsequent applications.*

*MycoApply Products are compatible with a variety of fungicides. Check our website or contact us for a full compatibility list before applying a fungicide.*